

Genomic and phenotypic characterization of a nonpigmented variant of *Pantoea vagans* biocontrol strain C9-1 lacking the 530-kb megaplasmid pPag3

Theo H.M. Smits¹, Fabio Rezzonico¹, Cosima Pelludat¹, Alexander Goesmann², Jürg E. Frey¹ & Brion Duffy¹

¹Division of Plant Protection, Agroscope Changins-Wädenswil ACW, Wädenswil, Switzerland; and ²CeBiTec, University of Bielefeld, Bielefeld, Germany

Correspondence: Theo H.M. Smits, Division of Plant Protection, Agroscope Changins-Wädenswil ACW, Schloss, Postfach, CH-8820 Wädenswil, Switzerland. Tel.: +41 44 783 6111; fax: +41 44 783 6305; e-mail: theo.smits@acw.admin.ch

Received 29 January 2010; revised 7 April 2010; accepted 7 April 2010.
Final version published online 10 May 2010.

DOI:10.1111/j.1574-6968.2010.01994.x

Editor: David Studholme

Keywords

Pantoea vagans; biological control; carotenoids; ampicillin; autoinduction; desferrioxamine.

Introduction

Pantoea is a diverse genus, with most species considered to be ubiquitous plant epiphytes and often isolated from a wide range of other environmental habitats (e.g., soil, water, insect/animal gut and clinical samples). Some plant isolates have demonstrated strong beneficial activity as biological control agents for pre- and postharvest fungal and bacterial diseases (Braun-Kiewnick *et al.*, 2000; Bonaterra *et al.*, 2005; Francés *et al.*, 2006). The type species, *Pantoea agglomerans*, has undergone extensive taxonomic rearrangement emerging from the *Enterobacter agglomerans*–*Erwinia herbicola* complex (Ewing & Fife, 1972; Rezzonico *et al.*, 2009). Recently, several closely related species, such as *P. vagans*, have been newly described based on molecular analysis (e.g., multilocus sequence analysis, DNA–DNA hybridization) (Brady *et al.*, 2008, 2009; Rezzonico *et al.*, 2009).

Strain C9-1 is an important biocontrol agent (Ishimaru *et al.*, 1988; Johnson & Stockwell, 1998) that is registered in the United States and Canada as Blight Ban C9-1 (NuFarms America). It is one of the most effective commercial agents

Abstract

A 530-kb megaplasmid pPag3 contributing 10.8% of the total genome of *Pantoea vagans* biocontrol strain C9-1 was sequenced. A rare nonpigmented variant C9-1W was obtained and shown to have lost pPag3, but retained all other plasmids (pPag1, pPag2). Phenotypic characterization of the variant confirmed the function of several annotated genes that may influence ecological fitness and efficacy. Metabolic profiling revealed important plasmid-based carbon utilization phenotypes. Plasmid loss resulted in thiamine auxotrophy, absence of carotenoid pigmentation, desferrioxamine diffusible siderophore biosynthesis, inherent ampicillin resistance and expression of AI-1 quorum-sensing signaling. This confirmed the functional expression of the corresponding genes located on pPag3 in *P. vagans*.

against fire blight, a major threat to global pome fruit production caused by the related enterobacterium *Erwinia amylovora* (Johnson & Stockwell, 1998). Recently, C9-1 has been reassigned from *P. agglomerans* to the novel species *P. vagans* based on its *gyrB* sequence (Brady *et al.*, 2009; Rezzonico *et al.*, 2009).

Pantoea agglomerans and *P. vagans* isolates are generally considered nonpathogenic. Most *P. agglomerans* isolates lack virulence determinants such as type III secretion systems (T3SS), while some contain a T3SS described as a non-pathogenic type (Rezzonico *et al.*, 2009). The phytopathogenicity of subspecies *P. agglomerans* pv. *gypsophilae* and pv. *betae* can be attributed to recently acquired large plasmids that carry a pathogenic type of T3SS and other virulence determinants (Ezra *et al.*, 2000; Mor *et al.*, 2001; Guo *et al.*, 2002; Manulis & Barash, 2003; Nissan *et al.*, 2006). Virulence and ecological fitness genes in phytopathogenic *P. agglomerans* pathovars, *Pantoea stewartii* and *Pantoea ananatis* are regulated by an autoinducer-1 quorum-sensing system involving *N*-acyl-homoserine lactone (AHL) signals (von

Bodman *et al.*, 2003; Morohoshi *et al.*, 2007; Chalupowicz *et al.*, 2008, 2009).

Several *Pantoea* species are yellow pigmented (Grimont & Grimont, 2005) due to production of carotenoids (Sandmann *et al.*, 1990; Hundle *et al.*, 1994). Nonpigmented variants have been reported, arising spontaneously at a low frequency (10^{-2} – 10^{-3}) after extended cultivation on nutrient-rich laboratory media (Chatterjee & Gibbins, 1971; Gantotti & Beer, 1982; Lindh *et al.*, 1991). These reports described physiological changes, such as thiamine deficiency and negative reactions with citrate or maltose, and lack of reversion to a wild-type phenotype, suggesting that such phenotypic changes are due to plasmid loss, although this has never been confirmed experimentally.

We have found that *P. vagans* C9-1 carries three plasmids: two plasmids of 168 kb (pPag1) and 166 kb (pPag2), and the 530-kb megaplasmid named pPag3 (Smits *et al.*, 2009). The phenotypic effects of these plasmids in *P. vagans* C9-1 have not been described previously. Sequence analysis of the megaplasmid revealed that carotenoid biosynthesis is encoded on plasmid pPag3. We obtained a nonpigmented variant of *P. vagans* C9-1 that lost the ability to synthesize thiamine and metabolize maltose, features that were also encoded by plasmid pPag3 genes. The aim of this study was to use the nonpigmented variant that lacks pPag3, representing over 10% of the total genome, in order to confirm functional phenotypes for annotated plasmid genes.

Materials and methods

Strains, growth media and conditions

Bacteria were routinely grown at 28 °C on Luria–Bertani (LB) (Sambrook *et al.*, 1989). Carbon source (glucose, sorbitol, maltose or sucrose) and thiamine ($5 \mu\text{g mL}^{-1}$) utilization assays were conducted in amended M9 minimal medium (Sambrook *et al.*, 1989). Resistance to ampicillin (2.5 – $200 \mu\text{g mL}^{-1}$) or tellurite ($50 \mu\text{g mL}^{-1}$) was determined on amended LB agar. Siderophore biosynthesis was assessed on chrome azurol S (CAS) agar (Schwyn & Neilands, 1987) with $5 \mu\text{L}$ of an LB preculture spotted onto the medium. The AHL biosensor strain *Agrobacterium tumefaciens* NTL4[pZLR4] was grown in LB medium containing $30 \mu\text{g mL}^{-1}$ gentamicin (Luo *et al.*, 2003).

Plasmid curing

The method of Gantotti & Beer (1982) was used to generate a nonpigmented variant of *P. vagans* C9-1. An LB culture of C9-1 wild type was incubated at 38 °C for 24 h, and 10^{-5} – 10^{-6} dilutions were plated onto LB agar and incubated at 37 °C for 5 days. The nonpigmented variant C9-1W that was obtained was tested for the presence of the three C9-1 plasmids using PCR. Oligonucleotides (Supporting Infor-

mation, Table S2) were synthesized by Sigma-Genosys (Steinheim, Germany). The HotStarTaq Master Mix kit (Qiagen, Hilden, Germany) was used as described by the supplier. Chromosomal DNA was isolated using the Wizard Genomic DNA Purification Kit (Promega, Madison, WI). PCR was performed as described elsewhere (Innis *et al.*, 1990). PCR products were visualized on 1.5% agarose gels (Sambrook *et al.*, 1989).

Metabolic profiling

The metabolic profiles of *P. vagans* C9-1 and C9-1W were obtained using Biolog GN2 and AN plates (Hayward, CA). Precultures were grown in M9 medium with 5 mM glucose and allowed to grow to the late stationary phase to ensure complete substrate utilization. Cultures were centrifuged at 4000 g and the cell pellets were washed once before resuspending in a fresh M9 medium. The attenuation at 600 nm ($A_{600\text{nm}}$) was set to 0.15 and each microtiter plate well was inoculated with 100 μL of this bacterial suspension. The plates were scored after 1, 2 and 5 days of incubation at 28 °C.

Autoinducer bioassay for the detection of AHL biosynthesis

For AHL bioassays, cell-free filtrates (150 μL) from stationary-phase cultures of *P. vagans* (16 h, 28 °C) were combined with 150 μL of a washed stationary-phase culture of *A. tumefaciens* NTL4[pZLR4] (Luo *et al.*, 2003) (16 h, 28 °C) in a fresh LB medium containing 0.1% 5-bromo-4-chloro-3-indolyl β -galactoside (X-Gal). The production of AHL was determined qualitatively by observing a change to blue in the color of the microculture over the course of 3 days.

Plasmid sequence analysis

The genome sequence of plasmid pPag3 from *P. vagans* C9-1 (Smits *et al.*, 2009) was annotated using GenDB (Meyer *et al.*, 2003) and was deposited at GenBank (Accession number CP001895). Sequence manipulations were performed using the Lasergene package (DNASTAR, Madison, WI). Additional BLAST searches (Altschul *et al.*, 1990) were performed at NCBI (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Results and discussion

The sequence of pPag3

The genome sequence of *P. vagans* C9-1 (Smits *et al.*, 2009) revealed a 530-kb plasmid, designated pPag3, encoding the carotenoid biosynthesis cluster *crtEXYIBZ* (Pvag_pPag30170–Pvag_pPag30175) as the most prominent feature. The encoded proteins share 91–97% sequence identity to the respective proteins of *P. agglomerans* pv. *milletiae* Wist 801 (GenBank: AB076662) and 70–89% to those of

P. ananatis 20D3^T (Misawa *et al.*, 1990). The plasmid also carries four thiamine biosynthesis genes (*thiOSGF*; Pvag_pPag30158–Pvag_pPag30161) and a complete maltose metabolism gene cluster (Pvag_pPag30206–Pvag_pPag30215). The genes have an identical order as in other Enterobacteria (i.e., *Escherichia coli*) (Blattner *et al.*, 1997; Dippel & Boos, 2005).

Based on the sequence annotation, the genetic information encoded on pPag3 corresponds with the previously described phenotypic characteristics of nonpigmented variants of *P. agglomerans* (i.e., thiamine deficiency, lack of maltose utilization, no pigmentation) (Chatterjee & Gibbins, 1971; Gantotti & Beer, 1982; Lindh *et al.*, 1991), indicating that the plasmids in both species likely have similar features. In addition, when multiplying the size of pPag3 (530 kb) with the average molecular weight of a base pair (660 Da), the molecular weight of pPag3 obtained is in agreement with the observed 350 MDa plasmid reported from *P. agglomerans* (ex *E. herbicola*) Eh112Y (Gantotti & Beer, 1982).

Curing of plasmid pPag3 and characterization of the variant

A nonpigmented variant of *P. vagans* C9-1, designated C9-1W, was obtained (Fig. 1). The identity of C9-1W as a derivative rather than as a contaminant was confirmed with 100% *gyrB* sequence identity compared with C9-1. The distinctive white colony color is attributable to loss of the carotenoid biosynthetic gene cluster located on pPag3. Genotyping with multiple primer pairs targeting parts of the three plasmids in *P. vagans* C9-1 confirmed the absence of pPag3 and the presence of pPag1 and pPag2.

The plasmid sequence data identified a thiamine biosynthetic cluster (*thiOSGF*). Whether these are required for thiamine biosynthesis was unclear because several other genes (*thiBCDEIJKLMPQ*) known from pathways in prokaryotes (Begley *et al.*, 1999; Settembre *et al.*, 2003) are found scattered on the *P. vagans* C9-1 chromosome (Smits




Fig. 1. *Pantoea vagans* C9-1 (left) and its nonpigmented variant C9-1W (right). The lack of yellow pigmentation could be attributed to the absence of the carotenoid biosynthesis gene cluster (*crtEXYIBZ*; Pvag_pPag30170–Pvag_pPag30175).

et al., 2009). When tested on glucose-amended minimal media, C9-1W only grew with a thiamine supplement. This confirms that plasmid-borne *thiOSGF* are essential for thiamine autotrophy in *P. vagans* C9-1, and may explain thiamine auxotrophy reported for the nonpigmented variant, plasmid-cured *P. agglomerans* (Chatterjee & Gibbins, 1971; Gantotti & Beer, 1982). Substitution of glucose with maltose in the minimal medium resulted in no growth regardless of the presence/absence of thiamine. This further demonstrates that maltose utilization (Dippel & Boos, 2005) is conferred by genes located on pPag3 (Pvag_pPag30206–Pvag_pPag30215). When *P. vagans* C9-1W was grown with sucrose or sorbitol as the sole carbon sources, thiamine was again found to be the critical parameter for growth. This confirms the thiamine auxotrophy of *P. vagans* C9-1W, and also the retention of pPag1 (containing sucrose metabolic genes) and pPag2 (containing sorbitol metabolic genes) in the variant.

Ampicillin resistance


Plasmid pPag3 also contains two genes with high sequence identity to a β -lactamase *bla* and its cognate regulator *ampR* (Pvag_pPag30395–Pvag_pPag30396). Ampicillin resistance has been reported to occur commonly in *P. agglomerans* clinical isolates (Cruz *et al.*, 2007). Unlike the wild-type strain *P. vagans* C9-1, variant C9-1W was sensitive to ampicillin at 200 $\mu\text{g mL}^{-1}$, suggesting that a functional β -lactamase is encoded on the plasmid.

Autoinducer biosynthesis

Autoinduction mediated by AHL signals has been well described in the plant pathogen *P. stewartii* ssp. *stewartii* (von Bodman *et al.*, 2003) and has been reported recently in the pathogens *P. agglomerans* pv. *gypsophilae* and *P. ananatis* (Morohoshi *et al.*, 2007; Chalupowicz *et al.*, 2008). Based on the sequence homology to the *pagRI* genes of *P. agglomerans* pv. *gypsophilae* (Chalupowicz *et al.*, 2008; Rezzonico *et al.*, 2009) the transcriptional regulator *pagR* and the AHL-synthase *pagI* genes (Pvag_pPag30141–Pvag_pPag30142) have been identified on plasmid pPag3.

Using an *A. tumefaciens* biosensor (Shaw *et al.*, 1997), AHL production was tested. For this purpose, the plant pathogenic strain *P. ananatis* LMG 2665 was added to the assay as a positive control. *Pantoea vagans* C9-1 has a positive autoinducer functional activity, but yields a weaker signal in the biosensor assay than *P. ananatis* LMG 2665 (Fig. 2). The variant *P. vagans* C9-1W lost this activity (Fig. 2), confirming that this strain is not able to produce detectable AHLs. Although the chromosome also contains a putative AHL synthase, located next to the *sdiA* gene encoding a LuxR-type transcriptional regulator (Lindsay & Ahmer, 2005; Smits *et al.*, 2009), it can be concluded from

Fig. 2. Detection of AHL production *Pantoea vagans* C9-1 and its nonpigmented variant C9-1W using the *Agrobacterium tumefaciens* biosensor assay. As a positive control, *Pantoea ananatis* LMG 2665^T was used; fresh LB medium was used as a negative control.



the results of the biosensor assay that this chromosomal gene is not involved in the synthesis of the AHLs that can be detected with the *A. tumefaciens* biosensor.

The PagRI quorum-sensing system plays a central role in the virulence of *P. agglomerans* pv. *gypsophilae* by regulating the expression of the T3SS (Chalupowicz *et al.*, 2009). Its role in the ecological behavior of *P. vagans* and *P. agglomerans* strains that have functional *pagRI* genes is currently unknown (Rezzonico *et al.*, 2009). The fact that most nonpathogenic strains lack a T3SS, however, suggests that *pagRI* may have additional non-virulence-related functions in phytopathogenic pathogens.

Iron acquisition

Siderophores are small molecules that bind Fe³⁺ with a high affinity and are synthesized by bacteria under iron starvation. The genome of *P. vagans* C9-1 contains biosynthetic genes for the catecholate siderophore enterobactin (*ent-fep*) and the hydroxamate siderophore desferrioxamine E (*dfo-JACS*), which were reported to be produced by the strain (Feistner & Ishimaru, 1996). Siderophore biosynthesis can be an important biocontrol trait, as the strain may be able to compete with phytopathogens for the already limited supply of iron in *planta*.

When spotted onto CAS siderophore indicator plates (Schwyn & Neilands, 1987), *P. vagans* C9-1 produces a large halo, indicative of siderophore synthesis, while variant C9-1W produces a small halo, just around the colony (Fig. 3). This difference can be attributed to the absence of the pPag3-encoded *dfoJACS* gene cluster (Pvag_pPag30339–Pvag_pPag30342), which confers the ability of desferrioxamine production to the strain.

Metabolic profiling using Biolog plates

Pantoea vagans C9-1W was compared with the wild-type strain *P. vagans* C9-1 for its substrate utilization pattern as determined by Biolog GN2 and AN plate assays (see for selected data Table S1). In general, growth with some of the compounds appeared to be slower than with the wild-type strain, and it cannot be excluded that this is influenced by the thiamine auxotrophy (thiamine was added for C9-1W to the minimal medium in the Biolog assays) or by the




Fig. 3. CAS agar plate with *Pantoea vagans* C9-1 (left) and its nonpigmented variant C9-1W (right). The lack of a large halo around C9-1W can be explained by the absence of the biosynthetic genes for desferrioxamine (*dfoJACS*; Pvag_pPag30339–Pvag_pPag30342). The weak signal indicates the production of chromosomally encoded enterobactin.

physiological differences in growth caused by the absence of pPag3.

Growth with maltose and maltotriose is abolished in *P. vagans* C9-1W due to the lack of the complete *mal* operon (Pvag_pPag30206–Pvag_pPag30215). Cellobiose, arbutin and salicin tested negative when using in C9-1W, but positive with the wild-type strain C9-1. These substrates are transported over the cytoplasmic membrane and channelled into the central pathways via a phosphotransferase system and a phosphohydrolase, respectively (An *et al.*, 2004, 2005). These functions are putatively encoded by two gene clusters on pPag3, *bgIBFG* (Pvag_pPag30318–Pvag_pPag30320) and *ascBFG* (Pvag_pPag30345–Pvag_pPag30437). The plasmid pPag3 contains the *gabTP* genes (Pvag_pPag30456–Pvag_pPag30457) (Niegemann *et al.*, 1993), described for their role in the uptake (GapP) and the initial transamination of γ -aminobutyrate (GABA) to succinate semialdehyde (GapT), which is subsequently channelled into the TCA cycle. Growth with GABA is retarded in C9-1W compared with the wild type, but not absent. Therefore, it is likely that there is an alternative pathway for growth with GABA in *P. vagans* C9-1W.

Growth with many organic acids is either retarded or absent in *P. vagans* C9-1W. This might partly be caused by the thiamine deficiency mentioned above. In addition, as plasmid pPag3 encodes several proteins involved in the uptake and conversion of organic acids, the lack of these functions may also contribute to these phenotypes in *P. vagans* C9-1W. The same may be true for the observed

Table 1. Presence of amplicons for pPag3 in *Pantoea vagans* strains

Amplicon	C9-1	LMG 24199 ^T	LMG 24196	LMG 24195
A	+	–	–	–
B	+	–	–	–
C	+	–	–	–
<i>pagRI</i>	+	+	+	+
D	+	+	+	+

Primer combinations and amplicon positions are listed in Table S2.

delay or the absence of growth with some of the amino acids, for which putative transporter- and conversion-encoding genes are also encoded on pPag3. However, as a direct link between annotated genes and a certain phenotype cannot be made based only on bioinformatic analysis, these observations remain hypothetical until further data are collected.

Presence and composition of pPag3 in other *P. vagans* strains

A spontaneous nonpigmented variant of *P. vagans* strain LMG 24196 was obtained on a rich medium plate under normal laboratory conditions. This variant was tested with the primers for *pagRI* (Rezzonico *et al.*, 2009) with no amplification, in contrast to a positive amplification in the wild-type parent LMG 24196 and the other two *P. vagans* strains (LMG 24195 and LMG 24199^T) (Brady *et al.*, 2009). This indicates that these autoinducer genes are also plasmid-borne in this strain.

Four PCR primer sets targeting pPag3 in genes encoding hypothetical proteins (amplicons A–C) and within the putative TonB-dependent siderophore receptor gene *fepA* (amplicon D) (Table 1) were used to screen the *P. vagans* strains. *Pantoea vagans* C9-1 was positive for all four amplicons, whereas strains LMG 24199^T, LMG 24195 and LMG 24196, were positive for amplicon D, but negative for amplicons A–C (Table 1). Amplicons A–C are within a 20 kb region on pPag3 (Table S2), suggesting that this part of the plasmid was acquired recently by *P. vagans* C9-1.

Conclusions

We have described here some phenotypic features for which the predicted genes are spread over the 530-kb plasmid pPag3 of *P. vagans* C9-1. This study confirms that plasmid loss can occur in *P. vagans* C9-1, albeit at a low frequency, even under conditions designed to obtain variants (e.g., rich media), as has been observed in *P. agglomerans* strains (Chatterjee & Gibbins, 1971; Gantotti & Beer, 1982; Lindh *et al.*, 1991). Several phenotypes that are lost along with the loss of plasmid pPag3 may be important for the ecological fitness of *P. vagans* C9-1, disfavoring the selection of non-pigmented variants in natural environments. Chief among

these are carotenoid pigmentation that can protect against environmental stresses (Dussault *et al.*, 2008; Jöhler *et al.*, 2010) and thiamine and siderophore biosynthesis that may improve competitiveness (Temple *et al.*, 2004; Dubuis *et al.*, 2006).

Acknowledgements

We thank V.O. Stockwell (Oregon State University, Corvallis, Oregon) for providing C9-1 genomic DNA and valuable discussions. We also thank T.A. Coutinho (FABI, University of Pretoria, South Africa) for the kind gift of the *P. vagans* LMG strains. This study was financed by the Swiss Federal Office for Agriculture (FOAG Fire Blight Control Project) and the Swiss State Secretariat for Education and Research (SBF C06.0069), conducted within the European Science Foundation research network COST Action 873.

References

- Altschul SF, Gish W, Miller W, Myers EW & Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- An CL, Lim WJ, Hong SY *et al.* (2004) Analysis of the *bgl* operon structure and characterization of b-glucosidase from *Pectobacterium carotovorum* subsp. *carotovorum* LY34. *Biosci Biotech Bioch* **68**: 2270–2278.
- An CL, Lim WJ, Hong SY, Shin EC, Kim MK, Lee JR, Park SR, Woo JG, Lim YP & Yun HD (2005) Structural and biochemical analysis of the *asc* operon encoding 6-phospho-b-glucosidase in *Pectobacterium carotovorum* subsp. *carotovorum* LY34. *Res Microbiol* **156**: 145–153.
- Begley TP, Downs DM, Ealick SE *et al.* (1999) Thiamin biosynthesis in prokaryotes. *Arch Microbiol* **171**: 293–300.
- Blattner FR, Plunkett G III, Bloch CA *et al.* (1997) The complete genome sequence of *Escherichia coli* K-12. *Science* **277**: 1453–1474.
- Bonaterre A, Camps J & Montesinos E (2005) Osmotically induced trehalose and glycine betaine accumulation improves tolerance to desiccation, survival and efficacy of the postharvest biocontrol agent *Pantoea agglomerans* EPS125. *FEMS Microbiol Lett* **250**: 1–8.
- Brady C, Cleenwerck I, Venter S, Vancanneyt M, Swings J & Coutinho T (2008) Phylogeny and identification of *Pantoea* species associated with plants, humans and the natural environment based on multilocus sequence analysis (MLSA). *Syst Appl Microbiol* **31**: 447–460.
- Brady CL, Venter SN, Cleenwerck I, Engelbeen K, Vancanneyt M, Swings J & Coutinho TA (2009) *Pantoea vagans* sp. nov., *Pantoea eucalypti* sp. nov., *Pantoea deleyi* sp. nov. and *Pantoea anthophila* sp. nov. *Int J Syst Evol Micr* **59**: 2339–2345.
- Braun-Kiewnick A, Jacobsen BJ & Sands DC (2000) Biological control of *Pseudomonas syringae* pv. *syringae*, the causative agent of basal kernel blight of barley, by antagonistic *Pantoea agglomerans*. *Phytopathology* **90**: 368–375.

- Chalupowicz L, Manulis-Sasson S, Itkin M, Sacher A, Sessa G & Barash I (2008) Quorum-sensing system affects gall development incited by *Pantoea agglomerans* pv. *gypsophylae*. *Mol Plant Microbe In* **21**: 1094–1105.
- Chalupowicz L, Barash I, Panijel M, Sessa G & Manulis-Sasson S (2009) Regulatory interactions between quorum-sensing, auxin, cytokinin, and the Hrp regulon in relation to gall formation and epiphytic fitness of *Pantoea agglomerans* pv. *gypsophylae*. *Mol Plant Microbe In* **22**: 849–856.
- Chatterjee AK & Gibbins LN (1971) Induction of nonpigmented variants of *Erwinia herbicola* by incubation at supraoptimal temperatures. *J Bacteriol* **105**: 107–112.
- Cruz AT, Cazacu AC & Allen CH (2007) *Pantoea agglomerans*, a plant pathogen causing human disease. *J Clin Microbiol* **45**: 1989–1992.
- Dippel R & Boos W (2005) The maltodextrin system of *Escherichia coli*: metabolism and transport. *J Bacteriol* **187**: 8322–8331.
- Dubuis C, Rolli J, Lutz M, Défago G & Haas D (2006) Thiamine-auxotrophic mutants of *Pseudomonas fluorescens* CHA0 are defective in cell-cell signaling and biocontrol factor expression. *Appl Environ Microb* **72**: 2606–2613.
- Dussault D, Caillet S, Le Tien C & Lacroix M (2008) Carotenoids' influence on radiotolerance of *Pantoea agglomerans*, a plant pathogen. *Lett Appl Microbiol* **47**: 208–213.
- Ewing WH & Fife MA (1972) *Enterobacter agglomerans* (Beijerinck) comb. nov. (the herbicola-lathyri bacteria). *Int J Syst Bacteriol* **22**: 4–11.
- Ezra D, Barash I, Valinsky L & Manulis S (2000) The dual function in virulence and host range restriction of a gene isolated from the pPATH_{Ehg} plasmid of *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant Microbe In* **13**: 683–692.
- Feistner GJ & Ishimaru C (1996) Proferrioxamine profiles of *Erwinia herbicola* and related bacteria. *Biometals* **9**: 337–344.
- Francés J, Bonaterra A, Moreno MC, Cabrefiga J, Badosa E & Montesinos E (2006) Pathogen aggressiveness and postharvest biocontrol efficiency in *Pantoea agglomerans*. *Postharvest Biol Tec* **39**: 299–307.
- Gantotti BV & Beer SV (1982) Plasmid-borne determinants of pigmentation and thiamine prototrophy in *Erwinia herbicola*. *J Bacteriol* **151**: 1627–1629.
- Grimont PAD & Grimont F (2005) Genus XXIII. *Pantoea*. Bergey's manual of systematic bacteriology. *The Proteobacteria, part B: the Gammaproteobacteria, Vol. 2*, 2nd edn (Brenner DJ, Krieg NR, Staley JT & Garrity GM, eds), pp. 713–720. Springer, New York.
- Guo M, Manulis S, Mor H & Barash I (2002) The presence of diverse IS elements and an *avrPphD* homologue that acts as a virulence factor on the pathogenicity plasmid of *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant Microbe In* **15**: 709–716.
- Hundle B, Alberti M, Nievelstein V, Beyer P, Kleinig H, Armstrong GA, Burke DH & Hearst JE (1994) Functional assignment of *Erwinia herbicola* Eho10 carotenoid genes expressed in *Escherichia coli*. *Mol Gen Genet* **245**: 406–416.
- Innis MA, Gelfand DH, Sninsky JJ & White TJ (1990) *PCR Protocols. A Guide to Methods and Applications*. Academic Press Inc., San Diego.
- Ishimaru CA, Klos EJ & Brubaker RR (1988) Multiple antibiotic production by *Erwinia herbicola*. *Phytopathology* **78**: 746–750.
- Johler S, Stephan R, Hartmann I, Kuehner KA & Lehner A (2010) Genes involved in yellow pigmentation of *Cronobacter sakazakii* ES5 and influence of pigmentation on persistence and growth under environmental stress. *Appl Environ Microb* **76**: 1053–1061.
- Johnson KB & Stockwell VO (1998) Management of fire blight: a case study in microbial ecology. *Annu Rev Phytopathol* **36**: 227–248.
- Lindh E, Kjaeldgaard P, Frederiksen W & Ursing J (1991) Phenotypical properties of *Enterobacter agglomerans* (*Pantoea agglomerans*) from human, animal and plant sources. *APMIS* **99**: 347–352.
- Lindsay A & Ahmer BMM (2005) Effect of *sdia* on biosensors of *N*-acylhomoserine lactones. *J Bacteriol* **187**: 5054–5058.
- Luo Z-Q, Su S & Farrand SK (2003) In situ activation of the quorum-sensing transcription factor TraR by cognate and noncognate acyl-homoserine lactone ligands: kinetics and consequences. *J Bacteriol* **185**: 5665–5672.
- Manulis S & Barash I (2003) *Pantoea agglomerans* pvs. *gypsophylae* and *betae*, recently evolved pathogens? *Mol Plant Pathol* **4**: 307–314.
- Meyer F, Goesmann A, McHardy AC *et al.* (2003) GenDB – an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res* **31**: 2187–2195.
- Misawa N, Nakagawa M, Kobayashi K, Yamano S, Izawa Y, Nakamura K & Harashima K (1990) Elucidation of the *Erwinia uredovora* carotenoid biosynthetic pathway by functional analysis of gene products expressed in *Escherichia coli*. *J Bacteriol* **172**: 6704–6712.
- Mor H, Manulis S, Zuck M, Nizan R, Coplin DL & Barash I (2001) Genetic organization of the *hrp* gene cluster and *dspAE/BF* operon in *Erwinia herbicola* pv. *gypsophylae*. *Mol Plant Microbe In* **14**: 431–436.
- Morohoshi T, Nakamura Y, Yamazaki G, Ishida A, Kato N & Ikeda T (2007) The plant pathogen *Pantoea ananatis* produces *N*-acylhomoserine lactone and causes center rot disease of onion by quorum sensing. *J Bacteriol* **189**: 8333–8338.
- Niegemann E, Schulz A & Bartsch K (1993) Molecular organization of the *Escherichia coli* *gab* cluster: nucleotide sequence of the structural genes *gabD* and *gabP* and expression of the GABA permease gene. *Arch Microbiol* **160**: 454–460.
- Nissan G, Manulis-Sasson S, Weinthal D, Mor H, Sessa G & Barash I (2006) The type III effectors HsvG and HsvB of gall-forming *Pantoea agglomerans* determine host specificity and function as transcriptional activators. *Mol Microbiol* **61**: 1118–1131.
- Rezzonico F, Smits THM, Montesinos E, Frey JE & Duffy B (2009) Genotypic comparison of *Pantoea agglomerans* plant and clinical strains to address biosafety and taxonomic issues. *BMC Microbiol* **9**: 204.

- Sambrook J, Fritsch EF & Maniatis T (1989) *Molecular Cloning: A laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory Press, New York.
- Sandmann G, Woods WS & Tuveson RW (1990) Identification of carotenoids in *Erwinia herbicola* and in a transformed *Escherichia coli* strain. *FEMS Microbiol Lett* **71**: 77–82.
- Schwyn B & Neilands JB (1987) Universal chemical assay for the detection and determination of siderophores. *Anal Biochem* **160**: 47–56.
- Settembre E, Begley TP & Ealick SE (2003) Structural biology of enzymes of the thiamin biosynthesis pathway. *Curr Opin Struct Biol* **13**: 739–747.
- Shaw PD, Ping G, Daly SL, Cha C, Cronan JE Jr, Rinehart KL & Farrand SK (1997) Detecting and characterizing *N*-acyl-homoserine lactone signal molecules by thin-layer chromatography. *P Natl Acad Sci USA* **94**: 6036–6041.
- Smits THM, Rezzonico F, Pelludat C, Stockwell VO, Goesmann A, Frey JE & Duffy B (2009) Complete genome sequencing of *Pantoea agglomerans* strain C9-1. *IOBC Bull* **43**: 375–378.
- Temple TN, Stockwell VO, Loper JE & Johnson KB (2004) Bioavailability of iron to *Pseudomonas fluorescens* strain A506 on flowers of pear and apple. *Phytopathology* **94**: 1286–1294.
- von Bodman SB, Dietz Bauer D & Coplin DL (2003) Quorum sensing in plant-pathogenic bacteria. *Annu Rev Phytopathol* **41**: 455–482.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Comparison of substrate spectrum between *P. vagans* C9-1 and the nonpigmented variant C9-1W using BIOLOG GN2 and AN plates.

Table S2. PCR primers used for gene amplification.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.